

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

Crop or Site	AI	Num. of Fields	% Base Acres Treated	Base Acres Treated	Cum. Acres Treated	Total Lbs AI
ALFALFA	FERROUS SULFATE	92	0.68	5,876	6,420	55
ALMOND	FERROUS SULFATE	15	0.12	875	1,355	58
BEAN, SUCCULENT	FERROUS SULFATE	2	0.61	162	162	2
CABBAGE	FERROUS SULFATE	2	0.13	34	128	1
CANTALOUPE	FERROUS SULFATE	5	0.92	380	380	6
CORN (FORAGE - FC	FERROUS SULFATE	4	0.02	72	72	1
COTTON	FERROUS SULFATE	126	0.99	8,152	11,462	84
EGGPLANT	FERROUS SULFATE	1	1.00	10	10	0
GRAPE	FERROUS SULFATE	4	0.03	96	96	2
GRAPE, WINE	FERROUS SULFATE	11	0.15	747	747	56
LEMON	FERROUS SULFATE	2	0.11	55	55	2
LETTUCE, HEAD	FERROUS SULFATE	4	0.11	185	500	3
LETTUCE, LEAF	FERROUS SULFATE	1	0.02	35	82	1
MELON	FERROUS SULFATE	3	0.40	84	84	1
NECTARINE	FERROUS SULFATE	5	0.22	80	80	4
N-OUTDR PLANTS IN	FERROUS SULFATE	1	0.24	80	80	1
ONION, DRY	FERROUS SULFATE	4	0.35	222	232	2
ORANGE	FERROUS SULFATE	5	0.03	55	55	8
PEACH	FERROUS SULFATE	4	0.03	30	30	2
PEPPER, FRUITING	FERROUS SULFATE	5	1.35	389	579	5
PLUM	FERROUS SULFATE	4	0.15	50	50	2
POMEGRANATE	FERROUS SULFATE	1	0.18	16	16	0
POTATO	FERROUS SULFATE	5	0.43	215	215	1
RICE	FERROUS SULFATE	1	0.03	150	150	0
SORGHUM/MILO	FERROUS SULFATE	2	2.46	138	138	0
TOMATO, PROCESS	FERROUS SULFATE	5	0.09	254	383	3
WALNUT	FERROUS SULFATE	6	0.06	123	123	5
WATERMELON	FERROUS SULFATE	4	0.83	159	159	1
WHEAT	FERROUS SULFATE	2	0.03	126	126	0
ARTICHOKE, GLOBE	IRON PHOSPHATE	7	1.42	357	570	110
BLACKBERRY	IRON PHOSPHATE	1	0.01	0	0	0
BRUSSELS SPROUT	IRON PHOSPHATE	1	0.00	0	0	0
CHERRY	IRON PHOSPHATE	3	0.29	80	80	2
GRAPE	IRON PHOSPHATE	1	0.01	47	47	1
GRAPE, WINE	IRON PHOSPHATE	4	0.02	111	111	31
GRAPEFRUIT	IRON PHOSPHATE	1	0.03	3	3	1
HERB, SPICE	IRON PHOSPHATE	1	0.04	1	2	0
LANDSCAPE MAINT	IRON PHOSPHATE	1	0.03	1	1	0
LETTUCE, LEAF	IRON PHOSPHATE	1	0.00	0	0	0
N-GRNHS FLOWER	IRON PHOSPHATE	5	0.63	17	18	5
N-GRNHS PLANTS IN	IRON PHOSPHATE	7	0.74	24	50	15
N-GRNHS TRANSPL	IRON PHOSPHATE	1	0.01	0	0	0
N-OUTDR FLOWER	IRON PHOSPHATE	4	0.59	73	73	25

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

Crop or Site	AI	Num. of Fields	% Base Acres Treated	Base Acres Treated	Cum. Acres Treated	Total Lbs AI
N-OUTDR PLANTS IN	IRON PHOSPHATE	32	1.57	523	1,003	243
N-OUTDR TRANSPL	IRON PHOSPHATE	1	0.16	18	18	4
ORANGE	IRON PHOSPHATE	3	0.03	55	55	0
PEPPER, FRUITING	IRON PHOSPHATE	1	0.00	0	0	0
STRAWBERRY	IRON PHOSPHATE	2	0.03	21	21	1
SWISS CHARD	IRON PHOSPHATE	1	0.00	0	0	0
TANGERINE	IRON PHOSPHATE	1	0.03	5	5	1
TOMATO	IRON PHOSPHATE	1	0.01	5	5	1
VEGETABLE	IRON PHOSPHATE	2	0.30	0	0	0
ALMOND	METALDEHYDE	1	0.00	2	2	13
ARTICHOKE, GLOBE	METALDEHYDE	25	4.36	1,097	1,892	1,504
AVOCADO	METALDEHYDE	54	1.57	838	851	433
BLACKBERRY	METALDEHYDE	1	1.46	11	11	18
BOK CHOY	METALDEHYDE	5	0.10	8	17	3
BROCCOLI	METALDEHYDE	1	0.03	51	51	51
BRUSSELS SPROUT	METALDEHYDE	4	0.92	33	33	37
CABBAGE	METALDEHYDE	1	0.02	5	5	42
CACTUS PEAR	METALDEHYDE	3	14.32	42	65	106
CHERRY	METALDEHYDE	2	0.17	47	64	25
CITRUS	METALDEHYDE	4	2.12	111	172	41
GRAPE, WINE	METALDEHYDE	9	0.22	1,110	1,111	463
GRAPEFRUIT	METALDEHYDE	25	2.86	261	669	197
LEMON	METALDEHYDE	259	24.67	12,637	31,881	14,138
LETTUCE, HEAD	METALDEHYDE	11	0.05	94	94	33
LETTUCE, LEAF	METALDEHYDE	1	0.00	1	1	0
LIME	METALDEHYDE	2	8.18	9	9	4
MINT	METALDEHYDE	2	0.09	5	8	6
N-GRNHS FLOWER	METALDEHYDE	25	6.77	188	339	301
N-GRNHS PLANTS IN	METALDEHYDE	91	11.93	390	998	1,836
N-GRNHS TRANSPL	METALDEHYDE	12	1.82	42	86	78
N-OUTDR FLOWER	METALDEHYDE	28	2.35	292	354	340
N-OUTDR PLANTS IN	METALDEHYDE	183	12.20	4,068	9,116	9,621
N-OUTDR TRANSPL	METALDEHYDE	19	1.19	134	152	157
ORANGE	METALDEHYDE	373	6.61	13,494	19,205	7,894
RASPBERRY	METALDEHYDE	12	3.25	129	129	142
STRAWBERRY	METALDEHYDE	51	1.71	1,208	1,420	999
TANGELO	METALDEHYDE	8	3.45	162	238	118
TANGERINE	METALDEHYDE	15	4.42	697	705	101
UNCULTIVATED AG	METALDEHYDE	1	0.00	6	6	6
UNCULTIVATED NOI	METALDEHYDE	1	0.01	2	2	2
WALNUT	METALDEHYDE	1	0.01	27	27	0
WATERCRESS	METALDEHYDE	1	6.40	10	10	0
GRAPE	METHIOCARB	1	0.01	29	29	5

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

Crop or Site	AI	Num. of Fields	% Base Acres Treated	Base Acres Treated	Cum. Acres Treated	Total Lbs AI
GRAPE, WINE	METHIOCARB	1	0.00	6	6	1
LEMON	METHIOCARB	2	0.17	87	211	29
N-GRNHS FLOWER	METHIOCARB	29	3.66	101	221	165
N-GRNHS PLANTS IN	METHIOCARB	57	6.17	202	247	598
N-GRNHS TRANSPL	METHIOCARB	2	0.16	4	4	12
N-OUTDR FLOWER	METHIOCARB	17	1.38	171	287	209
N-OUTDR PLANTS IN	METHIOCARB	79	3.35	1,117	1,849	1,634
N-OUTDR TRANSPL	METHIOCARB	2	0.07	8	16	13
ORANGE	METHIOCARB	2	0.07	134	134	9
UNCULTIVATED AG	METHIOCARB	1	0.03	60	60	90

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

AI	Lbs AI/ acre treated			Num. apps	Num. Applications per			
	Med rate	Min rate	Max rate		Med apps/ field	Min apps/ field	Max apps/ field	Num. WFE apps
FERROUS SULFATE	0.01	0.00	0.02	102	1.00	1.00	2.00	97.31
FERROUS SULFATE	0.04	0.01	0.09	19	1.00	1.00	1.00	15.87
FERROUS SULFATE	0.01	0.01	0.01	2	1.00	1.00	1.00	2.00
FERROUS SULFATE	0.01	0.01	0.01	7	3.50	3.00	4.00	7.00
FERROUS SULFATE	0.01	0.01	0.02	5	1.00	1.00	1.00	5.00
FERROUS SULFATE	0.01	0.01	0.01	4	1.00	1.00	1.00	0.71
FERROUS SULFATE	0.01	0.00	0.02	191	1.00	1.00	3.00	171.77
FERROUS SULFATE	0.01	0.01	0.01	1	1.00	1.00	1.00	1.0
FERROUS SULFATE	0.02	0.01	0.03	4	1.00	1.00	1.00	4.00
FERROUS SULFATE	0.14	0.03	0.14	11	1.00	1.00	1.00	10.56
FERROUS SULFATE	0.04	0.02	0.07	2	1.00	1.00	1.00	1.19
FERROUS SULFATE	0.01	0.01	0.01	17	4.50	4.00	5.00	10.2
FERROUS SULFATE	0.01	0.01	0.01	3	3.00	3.00	3.00	2.34
FERROUS SULFATE	0.02	0.01	0.02	5	2.00	2.00	2.00	1.91
FERROUS SULFATE	0.07	0.01	0.10	5	1.00	1.00	1.00	4.62
FERROUS SULFATE	0.02	0.02	0.02	1	1.00	1.00	1.00	1.00
FERROUS SULFATE	0.01	0.01	0.01	5	1.00	1.00	1.00	3.97
FERROUS SULFATE	0.14	0.14	0.14	5	1.00	1.00	1.00	3.54
FERROUS SULFATE	0.05	0.04	0.07	4	1.00	1.00	1.00	2.65
FERROUS SULFATE	0.01	0.01	0.02	8	2.00	1.00	2.00	8.00
FERROUS SULFATE	0.03	0.03	0.03	4	1.00	1.00	1.00	4.00
FERROUS SULFATE	0.02	0.02	0.02	1	1.00	1.00	1.00	1.0
FERROUS SULFATE	0.00	0.00	0.01	5	1.00	1.00	1.00	3.01
FERROUS SULFATE	0.00	0.00	0.00	1	1.00	1.00	1.00	0.58
FERROUS SULFATE	0.00	0.00	0.00	2	1.00	1.00	1.00	2.00
FERROUS SULFATE	0.01	0.01	0.01	8	2.00	1.00	2.00	5.90
FERROUS SULFATE	0.01	0.01	0.07	6	1.00	1.00	1.00	5.01
FERROUS SULFATE	0.01	0.01	0.01	4	1.00	1.00	1.00	3.79
FERROUS SULFATE	0.00	0.00	0.00	2	1.00	1.00	1.00	2.00
IRON PHOSPHATE	0.19	0.11	0.20	16	2.00	1.00	3.00	17.13
IRON PHOSPHATE	0.36	0.36	0.36	1	1.00	1.00	1.00	1.00
IRON PHOSPHATE	0.15	0.15	0.15	3	3.00	3.00	3.00	3.00
IRON PHOSPHATE	0.03	0.02	0.04	3	1.00	1.00	1.00	2.91
IRON PHOSPHATE	0.01	0.01	0.01	1	1.00	1.00	1.00	1.00
IRON PHOSPHATE	0.24	0.24	0.24	5	1.00	1.00	1.00	0.54
IRON PHOSPHATE	0.17	0.17	0.17	1	1.00	1.00	1.00	0.92
IRON PHOSPHATE	0.18	0.10	0.20	6	6.00	6.00	6.00	3.40
IRON PHOSPHATE	0.03	0.03	0.03	1	1.00	1.00	1.00	0.02
IRON PHOSPHATE	0.16	0.10	0.23	4	4.00	4.00	4.00	3.50
IRON PHOSPHATE	0.44	0.07	0.44	17	1.00	1.00	2.00	4.04
IRON PHOSPHATE	0.21	0.16	1.09	19	2.00	1.00	4.00	7.54
IRON PHOSPHATE	0.69	0.69	0.69	1	1.00	1.00	1.00	0.10
IRON PHOSPHATE	0.40	0.25	0.40	7	1.00	1.00	1.00	1.04

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

AI	Lbs AI/ acre treated			Num. apps	Num. Applications per			
	Med rate	Min rate	Max rate		Med apps/ field	Min apps/ field	Max apps/ field	Num. WFE apps
IRON PHOSPHATE	0.25	0.03	0.50	104	2.00	1.00	11.00	26.25
IRON PHOSPHATE	0.24	0.21	0.26	15	15.00	15.00	15.00	0.37
IRON PHOSPHATE	0.01	0.01	0.01	3	1.00	1.00	1.00	2.67
IRON PHOSPHATE	0.15	0.15	0.15	3	3.00	3.00	3.00	3.00
IRON PHOSPHATE	0.40	0.40	0.40	3	1.50	1.00	2.00	1.38
IRON PHOSPHATE	0.15	0.15	0.15	3	3.00	3.00	3.00	3.00
IRON PHOSPHATE	0.20	0.20	0.20	1	1.00	1.00	1.00	1.00
IRON PHOSPHATE	0.25	0.25	0.25	1	1.00	1.00	1.00	0.50
IRON PHOSPHATE	0.20	0.10	0.30	2	1.00	1.00	1.00	0.30
METALDEHYDE	8.75	8.75	8.75	1	1.00	1.00	1.00	0.03
METALDEHYDE	0.61	0.01	1.56	49	1.00	1.00	5.00	43.13
METALDEHYDE	0.63	0.01	2.00	70	1.00	1.00	2.00	33.21
METALDEHYDE	1.60	1.60	1.60	1	1.00	1.00	1.00	0.58
METALDEHYDE	0.07	0.02	0.18	12	2.00	1.00	2.00	6.72
METALDEHYDE	1.00	1.00	1.00	1	1.00	1.00	1.00	0.25
METALDEHYDE	1.00	1.00	1.00	4	1.00	1.00	1.00	1.80
METALDEHYDE	1.00	1.00	1.00	1	1.00	1.00	1.00	1.00
METALDEHYDE	1.41	1.41	1.41	3	1.00	1.00	1.00	4.36
METALDEHYDE	0.35	0.26	0.44	3	1.50	1.00	2.00	2.49
METALDEHYDE	0.40	0.25	0.40	5	1.00	1.00	1.00	4.28
METALDEHYDE	0.40	0.01	0.60	9	1.00	1.00	1.00	6.11
METALDEHYDE	0.80	0.10	1.33	40	1.00	1.00	3.00	28.27
METALDEHYDE	0.43	0.03	1.70	729	2.00	1.00	9.00	538.03
METALDEHYDE	0.40	0.25	0.40	11	1.00	1.00	1.00	9.53
METALDEHYDE	0.40	0.40	0.40	1	1.00	1.00	1.00	0.02
METALDEHYDE	0.42	0.04	0.80	2	1.00	1.00	1.00	2.00
METALDEHYDE	0.67	0.50	0.67	5	2.50	2.00	3.00	3.75
METALDEHYDE	1.50	0.20	5.45	133	3.00	1.00	17.00	45.12
METALDEHYDE	1.50	0.08	11.54	462	3.00	1.00	16.00	174.12
METALDEHYDE	0.58	0.03	6.97	67	3.00	1.00	11.00	20.13
METALDEHYDE	0.87	0.04	3.20	111	2.50	1.00	11.00	17.89
METALDEHYDE	1.28	0.10	7.00	1,015	3.00	1.00	20.00	243.0
METALDEHYDE	0.56	0.04	3.45	53	2.00	1.00	5.00	23.17
METALDEHYDE	0.34	0.02	1.42	602	1.00	1.00	4.00	489.5
METALDEHYDE	0.35	0.02	1.60	13	1.00	1.00	1.00	8.40
METALDEHYDE	0.80	0.00	1.75	65	1.00	1.00	2.00	46.62
METALDEHYDE	0.40	0.05	0.83	12	1.00	1.00	2.00	11.53
METALDEHYDE	0.08	0.08	1.00	23	1.00	1.00	3.00	9.45
METALDEHYDE	0.55	0.10	1.00	1	1.00	1.00	1.00	0.60
METALDEHYDE	0.94	0.94	0.94	1	1.00	1.00	1.00	1.00
METALDEHYDE	0.02	0.02	0.02	1	1.00	1.00	1.00	0.90
METALDEHYDE	0.01	0.01	0.01	1	1.00	1.00	1.00	1.00
METHIOCARB	0.18	0.18	0.18	1	1.00	1.00	1.00	0.97

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

AI	Lbs AI/ acre treated			Num. apps	Num. Applications per			
	Med rate	Min rate	Max rate		Med apps/ field	Min apps/ field	Max apps/ field	Num. WFE apps
METHIOCARB	0.19	0.19	0.19	1	1.00	1.00	1.00	0.07
METHIOCARB	0.13	0.13	0.14	4	2.00	1.00	3.00	3.91
METHIOCARB	0.66	0.03	3.00	115	3.00	1.00	8.00	44.21
METHIOCARB	1.63	0.01	29.43	246	2.00	1.00	16.00	47.83
METHIOCARB	3.27	0.57	4.49	12	6.00	3.00	9.00	0.65
METHIOCARB	0.75	0.01	1.75	67	3.00	1.00	8.00	20.66
METHIOCARB	1.50	0.02	16.34	296	2.00	1.00	13.00	31.44
METHIOCARB	0.82	0.08	4.67	17	8.50	2.00	15.00	3.55
METHIOCARB	0.10	0.05	0.14	2	1.00	1.00	1.00	1.47
METHIOCARB	1.50	1.50	1.50	1	1.00	1.00	1.00	0.86

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

	treated field		
	Med WFE/ field	Min WFE/ field	Max WFE/ field
AI			
FERROUS SULFATE	1.00	0.59	2.00
FERROUS SULFATE	1.00	0.08	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	3.50	3.00	4.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	0.19	0.17	0.21
FERROUS SULFATE	1.00	0.16	3.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	0.94	1.00
FERROUS SULFATE	0.60	0.19	1.00
FERROUS SULFATE	2.97	2.60	3.33
FERROUS SULFATE	2.34	2.34	2.34
FERROUS SULFATE	0.79	0.79	0.79
FERROUS SULFATE	0.98	0.90	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	0.72	0.50	1.00
FERROUS SULFATE	0.67	0.33	1.00
FERROUS SULFATE	2.00	1.00	2.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	0.78	0.13	1.00
FERROUS SULFATE	0.58	0.58	0.58
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	0.98	1.83
FERROUS SULFATE	0.97	0.75	1.00
FERROUS SULFATE	1.00	1.00	1.00
FERROUS SULFATE	1.00	1.00	1.00
IRON PHOSPHATE	1.83	1.00	3.00
IRON PHOSPHATE	1.00	1.00	1.00
IRON PHOSPHATE	3.00	3.00	3.00
IRON PHOSPHATE	1.00	1.00	1.00
IRON PHOSPHATE	1.00	1.00	1.00
IRON PHOSPHATE	0.14	0.08	0.20
IRON PHOSPHATE	0.92	0.92	0.92
IRON PHOSPHATE	3.40	3.40	3.40
IRON PHOSPHATE	0.02	0.02	0.02
IRON PHOSPHATE	3.50	3.50	3.50
IRON PHOSPHATE	0.50	0.50	1.00
IRON PHOSPHATE	0.49	0.23	1.18
IRON PHOSPHATE	0.10	0.10	0.10
IRON PHOSPHATE	0.26	0.23	0.29

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

	treated field		
	Med WFE/ field	Min WFE/ field	Max WFE/ field
AI			
IRON PHOSPHATE	0.21	0.01	4.00
IRON PHOSPHATE	0.37	0.37	0.37
IRON PHOSPHATE	1.00	1.00	1.00
IRON PHOSPHATE	3.00	3.00	3.00
IRON PHOSPHATE	0.69	0.63	0.75
IRON PHOSPHATE	3.00	3.00	3.00
IRON PHOSPHATE	1.00	1.00	1.00
IRON PHOSPHATE	0.50	0.50	0.50
IRON PHOSPHATE	0.15	0.10	0.20
METALDEHYDE	0.03	0.03	0.03
METALDEHYDE	1.00	0.29	6.58
METALDEHYDE	0.59	0.04	2.00
METALDEHYDE	0.58	0.58	0.58
METALDEHYDE	0.40	0.40	2.00
METALDEHYDE	0.25	0.25	0.25
METALDEHYDE	0.38	0.06	0.70
METALDEHYDE	1.00	1.00	1.00
METALDEHYDE	1.68	1.68	1.68
METALDEHYDE	1.24	1.00	1.49
METALDEHYDE	1.00	1.00	1.00
METALDEHYDE	0.76	0.51	0.98
METALDEHYDE	1.00	0.02	3.00
METALDEHYDE	1.71	0.08	8.57
METALDEHYDE	1.00	0.50	1.00
METALDEHYDE	0.02	0.02	0.02
METALDEHYDE	1.00	1.00	1.00
METALDEHYDE	1.88	1.75	2.00
METALDEHYDE	0.82	0.02	6.60
METALDEHYDE	0.92	0.07	10.69
METALDEHYDE	0.76	0.04	4.00
METALDEHYDE	0.39	0.02	1.38
METALDEHYDE	0.69	0.00	8.34
METALDEHYDE	0.72	0.25	3.00
METALDEHYDE	1.00	0.06	4.00
METALDEHYDE	0.95	0.11	1.00
METALDEHYDE	1.00	0.17	2.00
METALDEHYDE	1.00	1.00	2.00
METALDEHYDE	0.91	0.03	1.00
METALDEHYDE	0.60	0.60	0.60
METALDEHYDE	1.00	1.00	1.00
METALDEHYDE	0.90	0.90	0.90
METALDEHYDE	1.00	1.00	1.00
METHIOCARB	0.97	0.97	0.97

The production agricultural use of snail and slug pesticides on all crops in California in 2004. The measures of use are described in the accompanying table.

AI	treated field		
	Med WFE/ field	Min WFE/ field	Max WFE/ field
METHIOCARB	0.07	0.07	0.07
METHIOCARB	1.95	1.00	2.91
METHIOCARB	1.00	0.06	4.07
METHIOCARB	0.39	0.01	4.58
METHIOCARB	0.32	0.05	0.59
METHIOCARB	0.27	0.03	4.00
METHIOCARB	0.11	0.00	3.47
METHIOCARB	1.77	0.40	3.15
METHIOCARB	0.73	0.47	1.00
METHIOCARB	0.86	0.86	0.86